

## 2.5

# Play Ball!

## Absolute Value Equations and Inequalities

### LEARNING GOALS

In this lesson, you will:

- Understand and solve absolute values.
- Solve linear absolute value equations.
- Solve and graph linear absolute value inequalities on number lines.
- Graph linear absolute values and use the graph to determine solutions.

### KEY TERMS

- opposites
- absolute value
- linear absolute value equation
- linear absolute value inequality
- equivalent compound inequalities

All games and sports have specific rules and regulations. There are rules about how many points each score is worth, what is in-bounds and what is out-of-bounds, and what is considered a penalty. These rules are usually obvious to anyone who watches a game. However, some of the regulations are not so obvious. For example, the National Hockey League created a rule that states that a blade of a hockey stick cannot be more than three inches or less than two inches in width at any point. In the National Football League, teams that wear black shoes must wear black shoelaces and teams that wear white shoes must wear white laces. In the National Basketball Association, the rim of the basket must be a circle exactly 18 inches in diameter. Most sports even have rules about how large the numbers on a player's jersey can be!


Do you think all these rules and regulations are important? Does it really matter what color a player's shoelaces are? Why do you think professional sports have these rules, and how might the sport be different if these rules did not exist?

**PROBLEM 1** Opposites Attract? Absolutely!



1. Analyze each pair of numbers and the corresponding graph.


a. -2 and 2



b.  $-\frac{2}{3}$  and  $\frac{2}{3}$



c. -1.5 and 1.5



2. Describe the relationship between the two numbers.

3. What do you notice about the distance each point lies away from zero on each number line?

Two numbers that are an equal distance, but are in different directions, from zero on the number line are called **opposites**. The **absolute value** of a number is its distance from zero on the number line.



4. Write each absolute value.

a.  $|-2| = \underline{\hspace{2cm}}$

$|2| = \underline{\hspace{2cm}}$

b.  $|\frac{-2}{3}| = \underline{\hspace{2cm}}$

$|\frac{2}{3}| = \underline{\hspace{2cm}}$

c.  $|-1.5| = \underline{\hspace{2cm}}$

$|1.5| = \underline{\hspace{2cm}}$

5. What do you notice about each set of answers for Question 4?

How can you use each corresponding graph in Question 1 to verify your answers?



© 2012 Carnegie Learning

6. Determine the value of each. Show your work.

a.  $|3 - 8|$

b.  $|3| - |8|$

c.  $|4(5)|$


d.  $|-4| \cdot |5|$

e.  $|\frac{12}{-3}|$

f.  $\frac{|12|}{|-3|}$

2

7. Determine the solution(s) to each equation.



a.  $x = 5$

b.  $|x| = 5$

c.  $|x| = -5$

d.  $|x| = 0$

Use the number line as a tool to think about each solution.



8. Analyze each equation containing an absolute value symbol in Question 7. What does the form of the equation tell you about the possible number of solutions?

**PROBLEM 2** Too Heavy? Too Light? You're Out!

The official rules of baseball state that all baseballs used during professional games must be within a specified range of weights. The baseball manufacturer sets the target weight of the balls at 145.045 grams on its machines. The specified weight allows for a difference of 3.295 grams. This means the weight can be 3.295 grams greater than or less than the target weight.

**2**

1. Write an expression to represent the difference between a manufactured baseball's weight and the target weight. Use  $w$  to represent a manufactured baseball's weight.
  
2. Suppose the manufactured baseball has a weight that is greater than the target weight.
  - a. Write an equation to represent the greatest acceptable difference in the weight of a baseball.
  
  
  
  
  
  
  
  
  
  
  - b. Solve your equation to determine the greatest acceptable weight of a baseball.
  
3. Suppose the manufactured baseball has a weight that is less than the target weight.
  - a. Write an equation to represent the least acceptable difference in weight.



- b. Solve your equation to determine the least acceptable weight of a baseball.



The two equations you wrote can be represented by the **linear absolute value equation**  $|w - 145.045| = 3.295$ . In order to solve any absolute value equation, recall the definition of absolute value.



Consider this linear absolute value equation.

$$|a| = 6$$

There are two points that are 6 units away from zero on the number line: one to the right of zero, and one to the left of zero.

$$+(a) = 6 \quad \text{or} \quad -(a) = 6$$

$$a = 6 \quad \text{or} \quad a = -6$$

Now consider the case where  $a = x - 1$ .

$$|x - 1| = 6$$

If you know that  $|a| = 6$  can be written as two separate equations, you can rewrite any absolute value equation.

$$+(a) = 6 \quad \text{or} \quad -(a) = 6$$

$$+(x - 1) = 6 \quad \text{or} \quad -(x - 1) = 6$$

2



4. How do you know the expressions  $+(a)$  and  $-(a)$  represent opposite distances?



5. Determine the solution(s) to the linear absolute value equation  $|x - 1| = 6$ . Then check your answer.

$$+(x - 1) = 6$$

$$-(x - 1) = 6$$

The expressions  $+(x - 1)$  and  $-(x - 1)$  are opposites.



To solve each equation, would it be more efficient to distribute the negative or divide both sides of the equation by  $-1$  first?





6. Solve each linear absolute value equation. Show your work.

a.  $|x + 7| = 3$

b.  $|x - 9| = 12$

c.  $|3x + 7| = -8$

d.  $|2x + 3| = 0$

Before you start solving each equation, think about the number of solutions each equation may have. You may be able to save yourself some work—and time!



2

7. Cho, Steve, Artie, and Donald each solved the equation  $|x| - 4 = 5$ .

**Artie**

$$\begin{array}{l}
 |x| - 4 = 5 \\
 (x) - 4 = 5 \qquad -(x) - 4 = 5 \\
 (x) = 9 \qquad \qquad -x = 9 \\
 \qquad \qquad \qquad \qquad \qquad \qquad x = -9
 \end{array}$$

**Donald**

$$\begin{array}{l}
 |x| - 4 = 5 \\
 |x| = 9 \\
 (x) = 9 \qquad \qquad \qquad -(x) = 9 \\
 \qquad \qquad \qquad \qquad \qquad \qquad x = -9
 \end{array}$$

**Cho**

$$\begin{array}{l}
 |x| - 4 = 5 \\
 (x) - 4 = 5 \qquad -(x) - 4 = 5 \\
 x - 4 = 5 \qquad \qquad -x + 4 = 5 \\
 x = 9 \qquad \qquad \qquad -x = 1 \\
 \qquad \qquad \qquad \qquad \qquad \qquad x = -1
 \end{array}$$

**Steve**

$$\begin{array}{l}
 |x| - 4 = 5 \\
 (x) - 4 = +5 \qquad \qquad -(x) - 4 = -5 \\
 x = 9 \qquad \qquad \qquad -x - 4 = -5 \\
 \qquad \qquad \qquad \qquad \qquad \qquad -x = -1 \\
 \qquad \qquad \qquad \qquad \qquad \qquad x = 1
 \end{array}$$

© 2012 Carnegie Learning

- a. Explain how Cho and Steve incorrectly rewrote the absolute value equation as two separate equations.

- b. Explain the difference in the strategies that Artie and Donald used. Which strategy do you prefer? Why?

2

8. Solve each linear absolute value equation.

a.  $|x| + 16 = 32$

Consider isolating the absolute value part of the equation before you rewrite as two equations.

b.  $23 = |x - 8| + 6$

c.  $3|x - 2| = 12$



d.  $35 = 5|x + 6| - 10$



**PROBLEM 3** Too Big? Too Small? Just Right.

In *Too Heavy? Too Light? You're Out!* you determined the linear absolute value equation to identify the most and least a baseball could weigh and still be within the specifications. The manufacturer wants to determine all of the acceptable weights that the baseball could be and still fit within the specifications. You can write a **linear absolute value inequality** to represent this problem situation.

**2**

1. Write a linear absolute value inequality to represent all baseball weights that are within the specifications.
  - a. A manufactured baseball weighs 147 grams.
  - b. A manufactured baseball weighs 140.8 grams.
  - c. A manufactured baseball weighs 148.34 grams.




- d. A manufactured baseball weighs 141.75 grams.

2

3. Complete the inequality to describe all the acceptable weights, where  $w$  is the baseball's weight. Then use the number line to graph this inequality.


$$\underline{\hspace{2cm}} \leq w \leq \underline{\hspace{2cm}}$$



4. Raymond has the job of disposing of all baseballs that are not within the acceptable weight limits.
- a. Write an inequality to represent the weights of baseballs that Raymond can dispose of.



- b. Graph the inequality on the number line.






In Little League Baseball, the diameter of the ball is slightly smaller than that of a professional baseball.

5. The same manufacturer also makes Little League baseballs. For these baseballs, the manufacturer sets the target diameter to be 7.47 centimeters. The specified diameter allows for a difference of 1.27 centimeters.
- a. Denise measures the diameter of the Little League baseballs as they are being made. Complete the table to determine each difference. Then write the linear absolute value expression used to determine the diameter differences.

2

	Independent Quantity	Dependent Quantity
	Diameter of the Little League Baseballs	Target and Actual Diameter Difference
Units		
	6.54	
	8.75	
	7.39	
	5.99	
	8	
	9.34	
	7.47	
	$d$	

- b. Graph the linear absolute value function,  $f(d)$ , on a graphing calculator. Sketch the graph on the coordinate plane.



2

6. Determine the diameters of all Little League baseballs that fit within the specifications.
- a. Use your graph to estimate the diameters of all the Little League baseballs that fit within the specifications. Explain how you determined your answer.

- b. Algebraically determine the diameters of all the baseballs that fit within the specification. Write your answer as an inequality.

7. The manufacturer knows that the closer the diameter of the baseball is to the target, the more likely it is to be sold. The manufacturer decides to only keep the baseballs that are less than 0.75 centimeter from the target diameter.
  - a. Algebraically determine which baseballs will not fall within the new specified limits and will not be kept. Write your answer as an inequality.

2



- b. How can you use your graph to determine if you are correct?

### Talk the Talk





Absolute value inequalities can take four different forms as shown in the table. To solve a linear absolute value inequality, you must first write it as an **equivalent compound inequality**.



Absolute Value Inequality	Equivalent Compound Inequality
$ ax + b  < c$	$-c < ax + b < c$
$ ax + b  \leq c$	$-c \leq ax + b \leq c$
$ ax + b  > c$	$ax + b < -c$ or $ax + b > c$
$ ax + b  \geq c$	$ax + b \leq -c$ or $ax + b \geq c$




1. Solve the linear absolute value inequality by rewriting it as an equivalent compound inequality. Then graph your solution on the number line.
  - a.  $|x + 3| < 4$




© 2012 Carnegie Learning

b.  $6 \leq |2x - 4|$




2


c.  $|-5x + 8| + 2 < 25$



d.  $|x + 5| > -1$



e.  $|x + 5| < -1$



Be prepared to share your solutions and methods.